


# Food Additives & Contaminants

Simultaneous detection of DNA from 10 food allergens by ligation-dependent probe amplification <i>Alexandra Ehlert, Anja Demmel, Christine Hupfer, Ulrich Busch and Karl-Heinz Engel</i>	409
Monitoring of free glutamic acid in Malaysian processed foods, dishes and condiments <i>M. Khairunnisak, A.H. Azizah, S. Jinap and A. Nurul Izzah</i>	419
Comparative study of three screening tests, two microbiological tube tests, and a multi-sulphonamide ELISA kit for the detection of antimicrobial and sulphonamide residues in eggs <i>V. Gaudin, C. Hedou, A. Rault, P. Sanders and E. Verdon</i>	427
Comparison of a fluoroquinolone surface plasmon resonance biosensor screening assay with established methods <i>Stefan Weigel, Mariel G. Pikkemaat, J.W. Alexander Elferink, Patrick P.J. Mulder, Anne-Catherine Huet, Philippe Delahaut, Sonja Schittko, Ruth Flerus and Michel Nielen</i>	441
Assessment of endogenous androgen levels in meat, liver and testis of Iranian native cross-breed male sheep and bull by gas chromatography-mass spectrometry <i>R. Ahmadkhaniha, F. Kobarfard, N. Rastkari, M.R. Khoshayand, M. Amimi and A. Shafiee</i>	453
Use of glyceroltriheptanoate as marker for processed animal by-products: development and validation of an analytical method <i>C. von Holst, A. Boix, S. Bellorini, F. Serano, S. Androni, B. Verkuylen and R. Margry</i>	466
Residues of rotenone, azadirachtin, pyrethrins and copper used to control <i>Bactrocera oleae</i> (Gmel.) in organic olives and oil <i>V. Simeone, N. Baser, D. Perrelli, G. Cesari, H. El Bilali and P. Natale</i>	475
An isotope dilution headspace method with gas chromatography-mass spectrometry for determination of propylene oxide in food <i>Xu-Liang Cao and Jeannette Corriveau</i>	482
Impact of durum wheat milling on deoxynivalenol distribution in the outcoming fractions <i>G. Ríos, N. Zakhia-Rozis, M. Chaourand, F. Richard-Forget, M.F. Samson, J. Abecassis and V. Lullien-Pellerin</i>	487
<i>Fusarium</i> mycotoxin content of UK organic and conventional wheat <i>S.G. Edwards</i>	496
Occurrence of deoxynivalenol and its 3- $\beta$ -D-glucoside in wheat and maize <i>F. Berthiller, C. Dallasta, R. Corradini, R. Marchelli, M. Sulyok, R. Krska, G. Adam and R. Schuhmacher</i>	507
Reduction of fumonisin B <sub>1</sub> in extruded corn breakfast cereals with salt, malt and sugar in their formulation <i>M. Castells, A.J. Ramos, V. Sanchis and S. Marín</i>	512
Structural analysis of enniatin H, I, and MK1688 and beauvericin by liquid chromatography-tandem mass spectrometry (LC-MS/MS) and their production by <i>Fusarium oxysporum</i> KFCC 11363P <i>H.-H. Song, H.-S. Lee, G.-P. Lee, S.-D. Ha and C. Lee</i>	518
Moulds and mycotoxins in rice from the Swedish retail market <i>E. Fredlund, A.-M. Thim, A. Gidlund, S. Brostedt, M. Nyberg and M. Olsen</i>	527
New hierarchical classification of food items for the assessment of exposure to packaging migrants: use of hub codes for different food groups <i>P. Northing, P.K.T. Oldring, L. Castle and P.A.S.S. Mason</i>	534
Occurrence of 2,2,4-trimethyl-1,3-pentanediol monoisobutyrate (Texanol <sup>®</sup> ) in foods packed in polystyrene and polypropylene cups <i>M. Kempf, S. Ramm, T. Feuerbach and P. Schreier</i>	563
Cell transformation activities of abietic acid and dehydroabietic acid: safety assessment of possible contaminants in paper and paperboard for food contact use <i>K. Ohmori and Y. Kawamura</i>	568
Use of overall migration methodology to test for food-contact substances with specific migration limits <i>E.L. Bradley, L. Castle, S.M. Jickells, K.A. Mountfort and W.A. Read</i>	574



1944-0049(2009)26:4;1-E



Taylor & Francis  
Taylor & Francis Group