

# Food Additives & Contaminants

Formation of styrene during the Maillard reaction is negligible <i>R. Stadler</i>	583
Determination of synthetic drugs used to adulterate botanical dietary supplements using QTRAP LC-MS/MS <i>Y. Chen, L. Zhao, F. Lu, Y. Yu, Y. Chai and Y. Wu</i>	595
Levels of preservatives (sulfite, sorbate and benzoate) in New Zealand foods and estimated dietary exposure <i>P. Cressey and S. Jones</i>	604
Development and single-laboratory validation of an HPLC method for the determination of cyclamate sweetener in foodstuffs <i>M.J. Scotter, L. Castle, D.P.T. Roberts, R. MacArthur, P.A. Brereton, S.K. Hasnip and N. Katz</i>	614
Core food of the French food supply: second Total Diet Study <i>V. Sirot, J.L. Vélatier, G. Calamassi-Tian, C. Dubuisson, C. Ménard, A. Dufour and J.C. Leblanc</i>	623
Presence and metabolism of endogenous androgenic-anabolic steroid hormones in meat-producing animals: a review <i>J. Scarth, C. Akre, L. van Ginkel, B. Le Bizec, H. De Brabander, W. Korth, J. Points, P. Téale and J. Kay</i>	640
Confirmatory method for the determination of various acetylgestagens in animal kidney fat using liquid chromatography-tandem mass spectrometry <i>Edward Malone, Geraldine Dowling, Chris Elliott, Glenn Kennedy and Liam Regan</i>	672
Occurrence of $\alpha$ - and $\beta$ -nortestosterone residues in the urine of injured male cattle <i>D. Glenn Kennedy, H. Desmond Shortt, Steven R.H. Crooks, Paul B. Young, Henrietta J. Price, Wesley G. Smyth and S. Armstrong Hewitt</i>	683
Influence of physiological status on residues of lipophilic xenobiotics in livestock <i>D.J. MacLachlan</i>	692
Application of enzyme-linked immunosorbent assay for quantification of the insecticides imidacloprid and thiamethoxam in honey samples <i>Huixin Ma, Yanjun Xu, Qing X. Li, Ting Xu, Xintong Wang and Ji Li</i>	713
Control strategies against grey mould ( <i>Botrytis cinerea</i> Pers.: Fr) and corresponding fungicide residues in grapes and wines <i>Patrick Edder, Didier Ortelli, Olivier Viret, Emmanuelle Cognard, Alexandre De Montmollin and Olivier Zali</i>	719
Effect of particle size and spectral sub-range within the UV-VIS-NIR range using diffuse reflectance spectra on multivariate models in evaluating the severity of fusariosis in ground wheat <i>G. Balcerowska, R. Siuda, J. Skrzypczak, A. Łukanowski and C. Sadowski</i>	726
<i>In vitro</i> and <i>in vivo</i> characterization of mycotoxin-binding additives used for animal feeds in Mexico <i>A. Marroquín-Cardona, Y. Deng, J.F. Taylor, C.T. Hallmark, N.M. Johnson and T.D. Phillips</i>	733
Fusarium mycotoxins in milling streams from the commercial milling of maize imported to the UK, and relevance to current legislation <i>K.A. Scudamore and S. Patel</i>	744
Quantitative analysis of patulin in apple juice by thin-layer chromatography using a charge coupled device detector <i>Juliane Elisa Welke, Michele Hoeltz, Horacio Alberto Dottori and Isa Beatriz Noll</i>	754
Off-flavours in wines through indirect transfer of volatile organic compounds (VOCs) from coatings <i>M.D. Fumi, M. Lambri and D.M. De Faveri</i>	759



1944-0049(2009)26:5;1-D



**Taylor & Francis**  
Taylor & Francis Group