

Food Additives & Contaminants

PART A:

CHEMISTRY • ANALYSIS • CONTROL • EXPOSURE & RISK ASSESSMENT



Taylor & Francis
Taylor & Francis Group

Contents

Detection of rabbit and hare processed material in compound feeds by TaqMan real-time PCR <i>N. Pegels, I. López-Calleja, T. García, R. Martín and I. González</i>	771
Determination of eight synthetic dyes in foodstuffs by green liquid chromatography <i>M. Hajimalmoodi, M. Afsharimanesh, G. Moghaddam, N. Sadeghi, M.R. Oveisi, B. Jannat, E. Pirhadi, F. Zamani Mazdeh and H. Kanan</i>	780
Monitoring the adulteration of milk with melamine: a visualised sensor array approach <i>L. Yang, D. Huo, Y. Jiang, C. Hou and S. Zhang</i>	786
Monitoring of ractopamine concentration in the mixture of this feed additive with vitamin mineral complex and with swine feed by HPLC <i>E.F. Freire, K.B. Borges, H. Tanimoto, R.T. Nogueira, L.C.T. Bertolini and C.M. de Gaitani</i>	796
Simultaneous determination of chloramphenicol, florfenicol and florfenicol amine in ham sausage with a hybrid chemiluminescent immunoassay <i>X. Tao, H. Jiang, X. Yu, J. Zhu, X. Wang, Z. Wang, L. Niu, X. Wu and J. Shen</i>	804
A simple method for the determination of fluoroquinolone residues in tilapia (<i>Oreochromis niloticus</i>) and pacu (<i>Piaractus mesopotamicus</i>) employing LC-MS/MS QToF <i>S.P. Quesada, J.A.R. Paschoal and F.G. Reyes</i>	813
Comparative study on disappearance trends of captan and trifloxystrobin residues on fruit and apple tree leaves using internal normalisation method <i>S. Sadlo, M. Duda, B. Piechowicz and A. Jaźwa</i>	826
Factors affecting the estimated probabilistic acute dietary exposure to captan from apple consumption <i>A. Zentai, J. Sali, I.J. Szabó, M. Szeitzné-Szabó, Á. Ambrus and A. Vársárhelyi</i>	833
Multiplex immunoassay for persistent organic pollutants in tilapia: comparison of imaging- and flow cytometry-based platforms using spectrally encoded paramagnetic microspheres <i>A. Meimaridou, W. Haasnoot, W.L. Shelver, M. Franek and M.W.F. Nielen</i>	843
Preparation and application of an immunoaffinity column based on an antibody with strong affinity and packing material with good stability <i>H. Zhang, M. Wang, Y. Wang, Q. Li, J. Zhou, F. Huo and B. Tang</i>	853
Effect of quercetin on the toxicokinetics of ochratoxin A in rats <i>Z. Abbas, R. Blank, S. Wein and S. Wolffram</i>	861
Effect of food processing on exposure assessment studies with mycotoxins <i>G. Cano-Sancho, V. Sanchez, A.J. Ramos and S. Marín</i>	867
Moniliformin analysis in maize samples from North-West Italy using multifunctional clean-up columns and the LC-MS/MS detection method <i>V. Scarpino, M. Blandino, M. Negre, A. Reyneri and F. Vanara</i>	876
Migration of mineral oil, photoinitiators and plasticisers from recycled paperboard into dry foods: a study under controlled conditions <i>M. Biedermann, J.-E. Ingenhoff, M. Zurfluh, L. Richter, T. Simat, A. Harling, W. Altkofer, R. Helling and K. Grob</i>	885
Migration of perfluoroalkyl acids from food packaging to food simulants <i>Y. Xu, G.O. Noonan and T.H. Begley</i>	899
Simulation of the migration of mineral oil from recycled paperboard into dry foods by Tenax® <i>M. Zurfluh, M. Biedermann and K. Grob</i>	909