

Volume 30 Number 8

August 2013

ISSN 1944-0049

Food Additives & Contaminants

PART A:

CHEMISTRY • ANALYSIS • CONTROL • EXPOSURE & RISK ASSESSMENT

Risk Control and Food Safety

**Proceedings of the 11th UJNR International Symposium on Toxic Microorganisms
held in Tokyo March 4–9, 2012**

Guest Editors: Shigeki Yamamoto and Kenneth A. Voss



Taylor & Francis
Taylor & Francis Group

Food Additives & Contaminants: Part A

Volume 30

Number 8

August 2013

Contents

Foreword

Risk control and food safety
S. Yamamoto and K.A. Voss

1349

Articles

Quantitative determination of paralytic shellfish toxins in cultured toxic algae by LC-MS/MS <i>R. Watanabe, R. Matsushima, T. Harada, H. Oikawa, M. Murata and T. Suzuki</i>	1351
LC-MS/MS analysis of palytoxin analogues in blue humphead parrotfish <i>Scarus ovifrons</i> causing human poisoning in Japan <i>T. Suzuki, R. Watanabe, R. Matsushima, K. Ishihara, H. Uchida, S. Kikutsugi, T. Harada, H. Nagai, M. Adachi, T. Yasumoto and M. Murata</i>	1358
<i>Kudoa septempunctata</i> was recognised by Toll-like receptor 2 produced by a RAW 264 macrophage-like cell line <i>T. Ohnishi, R. Oyama, H. Furusawa, N. Ohba, Y. Kamata and Y. Sugita-Konishi</i>	1365
Utility of the phylotoxigenic relationships among trichothecene-producing <i>Fusarium</i> species for predicting their mycotoxin-producing potential <i>M. Watanabe, T. Yonezawa, Y. Sugita-Konishi and Y. Kamata</i>	1370
Breeding aflatoxin-resistant maize lines using recent advances in technologies – a review <i>R.L. Brown, A. Menkir, Z.-Y. Chen, D. Bhatnagar, J. Yu, H. Yao and T.E. Cleveland</i>	1382
Exposure and risk assessment for ochratoxin A and fumonisins in Japan <i>Y. Sugita-Konishi, Y. Kamata, T. Sato, T. Yoshinari and S. Saito</i>	1392
Toxicological evaluation and bioaccumulation potential of lolitrem B, endophyte mycotoxin in Japanese black steers <i>N. Shimada, M. Yoshioka, O. Mikami, N. Tanimura, N. Yamanaka, M. Hanazumi, F. Kojima and S. Miyazaki</i>	1402
Detection of masked mycotoxins derived from type A trichothecenes in corn by high-resolution LC-Orbitrap mass spectrometer <i>H. Nakagawa, S. Sakamoto, Y. Sago, M. Kushiro and H. Nagashima</i>	1407
Alkaline cooking (nixtamalisation) and the reduction in the <i>in vivo</i> toxicity of fumonisin-contaminated corn in a rat feeding bioassay <i>K.A. Voss, R.T. Riley, N.D. Moore and T.D. Burns</i>	1415
Genomic paradigms for food-borne enteric pathogen analysis at the USFDA: case studies highlighting method utility, integration and resolution <i>C.A. Elkins, M.L. Kotewicz, S.A. Jackson, D.W. Lacher, G.S. Abu-Ali and I.R. Patel</i>	1422
Recent developments in molecular sub-typing of <i>Listeria monocytogenes</i> <i>A.R. Datta, P. Laksanalamai and M. Solomotis</i>	1437
Growth of <i>Listeria monocytogenes</i> in refrigerated ready-to-eat foods in Japan <i>Y. Okada, I. Ohnuki, H. Suzuki and S. Igimi</i>	1446
Prevalence of the main food-borne pathogens in retail food under the national food surveillance system in Japan <i>Y. Hara-Kudo, H. Konuma, Y. Kamata, M. Miyahara, K. Takatori, Y. Onoue, Y. Sugita-Konishi and T. Ohnishi</i>	1450
<i>Providencia alcalifaciens</i> causes barrier dysfunction and apoptosis in tissue cell culture: potent role of lipopolysaccharides on diarrheogenicity <i>H. Asakura, Y. Momose, C.-H. Ryu, F. Kasuga, S. Yamamoto, S. Kumagai and S. Igimi</i>	1459